



# WELCOME TO KINGS ONLINE PLANT DOCTOR

## CODLING MOTH

Cydia Pomonella

Codling Moth lay their eggs on pip fruit such as apples, pears and quince, as well as on walnuts.

Upon hatching the larvae burrows into the fruit and then feeds on the flesh and pips for approximately 3 weeks, before leaving the fruit to pupate, or overwinter in the ground.

The adult moths are 1 -2 cm long, and often hard to spot as their mottled wings blend well with tree bark. The larvae overwinter within silken cocoons in the soil, in leaf litter around the base of trees, or under loose scales of bark.

In Auckland the moths appear from September through till February, and during this period the moths mate. After mating females can lay between 30-70 tiny disc shaped eggs on the fruit, nuts, leaves and fruiting spurs.

### PREVENTION

Preventing the issue is always better than trying to cure it. As the Caterpillars pupate over winter and like to hide in various nooks and crannies, under bark, leaves, weeds and in infected fruit that are around trees, it is best to keep the area free of suitable places to pupate.

Give the trunk and branches a scrub (not too hard though) with a stiff brush to remove any loose bark that they may be hiding in.

### NATURAL TREATMENT

Spraying through spring with [BioNeem](#) may also help to smother eggs.

One remedy is to wrap the tree trunk in corrugated cardboard. The theory is that the caterpillar will find it an attractive hiding place and spin their cocoon in there. You can then remove and burn it before they hatch in spring.

To be most effective, place the cardboard as high up the trunk as possible (but below any fruit), grease or make a sticky band below the cardboard to stop them passing on by, and remove and burn frequently from early summer to after harvest.

### OTHER TREATMENT

When the trees bloom in spring, the adults will start to mate and lay eggs. This is the time to place a [Codling Moth Trap](#) in your trees. The trap uses a synthetic replica of the scent of the female to attract and trap the male moth, which reduces the chance of breeding and, therefore, fewer eggs. Replace the [pheromone](#) every 6-8 weeks.

One trap per 3-5 trees within a 15m radius is enough. Monitor the number of moths being caught, if there are more than 15 trapped moths in a 2 week period it is advisable to use additional traps or spray with an insecticide such as [Success Ultra](#).

When using sprays and chemicals always read the label and follow instructions carefully. Spray in the evening to avoid harming beneficial insects.



## CODLING MOTH PRODUCT RECOMMENDATIONS

Shop the Kings online store for all your gardening needs



### CODLING MOTH PHEROMONE TRAP

Natural protection against Codling Moth. Odourless poison-free weatherproof naturally safe for use around other beneficial insects.



### SUCCESS ULTRA INSECT CONTROL

The next generation solution in insect control. Success ULTRA helps keep your garden plants from being attacked by common caterpillars plus other insect pests like thrips, codling moth.



*Everything you need!*

FOR MORE ADVICE  
CONTACT US

0800 PLANTS  
09 410 9726

[www.plantdoctor.co.nz](http://www.plantdoctor.co.nz)  
[plantdoctor@kings.co.nz](mailto:plantdoctor@kings.co.nz)