



WELCOME TO KINGS ONLINE PLANT DOCTOR

GUAVA MOTH

Coscinoptycha improbana

A small moth, with a wingspan of approximately 15mm.

Originally from Australia, Guava Moth was first found in New Zealand in Northland in 1997.

The pest damages a wide range of fruit including feijoas, guavas, macadamia nuts, stone fruit and citrus. Female moths lay their eggs on the fruit, after hatching the larvae burrow into the flesh, causing significant damage and loss.

Guava moths are speckled black and white with a wingspan of approximately 15mm, while larvae are pink and can grow up to 8mm long. Guava moth can breed all year round

Signs to look out for include: premature fruit drop, exit holes or small discolorations, fruit may appear bruised, during the later stages the larvae's excrement may be visible. A pin-prick sized hole from the larvae burrowing in can sometimes be seen.

PREVENTION

When tackling the problem it's worth coordinating with neighbours to help avoid the build up of severe infestations on surrounding properties.

Prevent the moth from laying their eggs on the fruit by draping [Bug Net](#) over the tree and pinning it to the ground. Alternatively you can cut the net into sections and wrap around fruit.

Moths pupate in fallen fruit and leaf litter so keep the area under your trees free of grass, weeds and fallen fruit. Avoid composting infected fruit as the moths can complete their lifecycle in some compost systems.

Note: Wait till after flowering before covering the trees with [Bug Net](#) to allow pollination.

NATURAL TREATMENT

Another effective method of control is to use the [Little Bugga Guava Moth Trap](#). Unlike the pheromone trap it will catch female moths along with other species of moths, including codling moth.

In addition, spray with [GroSafe BioNeem](#) as fruit start to form.

Remove infected fruit and dispose of carefully to avoid spreading the larvae, avoid composting it on your property.

If you have chickens, fencing them in around your fruit trees will also help kill the grubs as they pupate in the soil.

OTHER TREATMENT

Current methods of control are somewhat limited. Use multiple methods for best results, and if possible work along with your neighbours for better control.

Monitor numbers or solve low-level infestations by placing [Guava Moth Traps](#) in your trees. One trap per 3 trees should be enough. The trap uses a pheromone that replicates the scent of the female moth, enticing the male moths into the trap and trapping them. Replace pheromone every 6 weeks with a [Easy Trap Guava Moth Refill](#).

If you are catching 5 or more moths over a two week period spray with [GroSafe BioNeem](#) twice a week on and around fruit until you stop catching the moths. Be especially vigilant when the flowers fall and the fruit first starts to form, as this is when the moths are laying their eggs.

When using sprays and chemicals always read the label and follow instructions carefully. Spray in the evening to avoid harming beneficial insects.



GUAVA MOTH PRODUCT RECOMMENDATIONS

Shop the Kings online store for all your gardening needs



GUAVA MOTH PHEROMONE TRAP

Naturally attracts the moth by using a pheromone lure then traps them before they damage your fruit.



BIONEEM INSECTICIDE

BioNeem is a broad spectrum insecticide derived from the Neem tree, for the control aphids, thrips and other insect pests.



Everything you need!

FOR MORE ADVICE
CONTACT US

0800 PLANTS
09 410 9726

www.plantdoctor.co.nz
plantdoctor@kings.co.nz